



## Secondary metabolites produced by fungi derived from a microbial mat encountered in an iron-rich natural spring

Alexandra L. Gereá<sup>a,†</sup>, Katie M. Branscum<sup>b,†</sup>, Jarrod B. King<sup>a</sup>, Jianlan You<sup>a</sup>, Douglas R. Powell<sup>b</sup>, Andrew N. Miller<sup>c</sup>, John R. Spear<sup>d</sup>, Robert H. Cichewicz<sup>a,\*</sup>

<sup>a</sup> Natural Products Discovery Group, Department of Chemistry and Biochemistry, Stephenson Life Sciences Research Center, 101 Stephenson Parkway, Room 1000, University of Oklahoma, Norman, OK 73019-5251, USA

<sup>b</sup> Department of Chemistry and Biochemistry, Stephenson Life Sciences Research Center, 101 Stephenson Parkway, Room 1000, University of Oklahoma, Norman, OK 73019-5251, USA

<sup>c</sup> University of Illinois, Illinois Natural History Survey, 1816 South Oak Street, Champaign, IL 61820-6970, USA

<sup>d</sup> Department of Civil and Environmental Engineering, Colorado School of Mines, Golden, CO 80401-1887, USA

### ARTICLE INFO

#### Article history:

Received 27 April 2012

Revised 29 May 2012

Accepted 30 May 2012

Available online 9 June 2012

#### Keywords:

Biofilm

Fungi

Microbial mat

Secondary metabolites

### ABSTRACT

A collection of fungal isolates was obtained from a complex microbial mat, which occupied an iron-rich freshwater spring that feeds into Clear Creek, Golden, Colorado, USA. Two of the fungal isolates, a Glomeromycete (possibly *Entrophospora* sp.) and a Dothideomycete (possibly *Phaeosphaeria* sp.), were investigated for bioactive secondary metabolites. In total, six new compounds consisting of clearanols A–E (**5**, **6**, **10–12**) and disulochrin (**7**) were purified and their structures were determined. Disulochrin exhibited modest antibacterial activity against methicillin-resistant *Staphylococcus aureus*, whereas clearanol C showed weak inhibitory activity against *Candida albicans* biofilm formation.

© 2012 Elsevier Ltd. All rights reserved.

Polymicrobial biofilms and mats are found in many natural settings at the interfaces between liquids and solid substrates.<sup>1–5</sup> Recognizing that these types of environments may provide ideal conditions for promoting interspecies metabolic exchange within and between all three domains of life,<sup>6</sup> we have sampled a limited number of naturally-occurring microbial mats for new secondary-metabolite-producing fungi. During the summer of 2010, one of the many springs that feed into Clear Creek near Golden, CO, USA (39°45'15.0" North, 105°13'39.9" West, elevation 1729 m) was host to a luxuriant microbial mat that appeared bright green due to an abundance of photosynthetic algae and cyanobacteria. However, by early autumn, the mat had developed an intense orange hue as iron-oxidizing *Gallionella*<sup>7</sup> became dominant in the iron-rich waters of the spring (4–7 mg/L iron in the spring versus 0.01–1.4 mg/L within the creek). Closer inspection of the mat revealed that a biodiverse mixture of other bacteria, as well as Archaea and Eukarya was present and a sample was collected for the purpose of procuring fungal isolates. In this Letter, we describe the structure assignments and bioassay screening results for metabolites purified from two fungal isolates obtained from the Clear

Creek microbial mat. The results from this investigation provide compelling evidence that naturally-occurring microbial communities offer unique opportunities for obtaining fungi that generate new bioactive compounds.

The Clear Creek mat sample was mechanically disrupted and aliquots of the microbial suspension were placed on Petri plates containing either Czapek or soil-extract agar with chloramphenicol. After two weeks of incubation at room temperature, fungal colonies were transferred to Czapek agar plates. This afforded 32 fungal isolates (20 from Czapek agar and 12 from soil-extract agar) that appeared distinct based on gross morphological features. Small-scale shake-flask cultures of the isolates were prepared and extracted with ethyl acetate. Upon removal of the solvent, the remaining organic residues were tested in our assay panel for antibacterial, antifungal, cancer cell cytotoxicity, and inhibition of *Candida* biofilm formation. The fungal isolates were also subjected to sequencing of their nuclear ribosomal internal transcribed spacer (ITS) regions. BLAST searches performed on the sequence data revealed that the majority of the fungi were taxonomically affiliated with isolates that were scantily studied or had not been subjected to prior analyses of their secondary metabolites (based on queries of the SciFinder<sup>®</sup> and Dictionary of Natural Products<sup>®</sup> databases). Two of the relatively uncommon isolates, a Glomeromycete (possibly *Entrophospora* sp., GenBank accession

\* Corresponding author. Tel.: +1 405 325 6969; fax: +1 405 325 6111.

E-mail address: [rhcichewicz@ou.edu](mailto:rhcichewicz@ou.edu) (R.H. Cichewicz).

† To be considered joint first authors.

no. JQ958304) and a Dothideomycete (possibly *Phaeosphaeria* sp., GenBank accession no. JQ958305), were chosen for scale-up chemical investigations.

An examination of different media formulations led to the observation that the Glomeromycete isolate grew particularly well under static conditions on a medium composed of Cheerios® breakfast cereal supplemented with a 0.3% sucrose solution. The fungal cultures were extracted with MeOH to give a dark reddish-rust colored extract that was subjected to a combination of silica gel chromatography and C18 HPLC. This provided several known compounds in substantial quantities including sulochrin (**1**) (218 mg), hydroxysulochrin (**2**) (12 mg), questinol (**3**) (57 mg), questin (**4**) (11 mg), and three new compounds (**5–7**) (Fig. 1).

Compound **5** (30 mg)<sup>8</sup> (Fig. 1) was assigned the molecular formula C<sub>11</sub>H<sub>14</sub>O<sub>5</sub> based on HRESIMS (*m/z* 227.0909 [M+H]<sup>+</sup>, calcd for C<sub>11</sub>H<sub>15</sub>O<sub>5</sub> 227.0914), which corresponded to five degrees of unsaturation. Inspection of the <sup>1</sup>H NMR, <sup>13</sup>C NMR, and <sup>1</sup>H–<sup>13</sup>C HSQC data provided evidence for five quaternary carbons, two methines, two methylenes, and two methyl groups. The <sup>1</sup>H–<sup>13</sup>C HMBC correlations (Fig. 2) were instrumental in piecing together compound **5** as the only reasonable planar structure for this metabolite. The structure of **5** was later confirmed by single crystal X-ray diffraction,<sup>9</sup> which also supported a 7*R* absolute configuration for the metabolite (Fig. 3) (determined by Hooft parameter, refer to the Supplementary data for additional details). Examination of the SciFinder database revealed that although the planar version of this compound had been previously assigned a CAS registry number (1235630-76-8), no references were available concerning its spectroscopic properties or synthetic/biosynthetic origins. We have given compound **5** the trivial name clearanol A.




Figure 1. Structures of clearanols A–E (**5**, **6**, **10–12**) and disulochrin (**7**).




Figure 2. Key <sup>2,3</sup>J<sub>H-C</sub> <sup>1</sup>H–<sup>13</sup>C HMBC correlations used to determine the structures of **5**, **6**, **10–12**.




Figure 3. Structure of **5** determined by single crystal X-ray diffraction experiment.

Compound **6** (0.7 mg)<sup>10</sup> (Fig. 1) exhibited <sup>1</sup>H and <sup>13</sup>C NMR data that were similar to those acquired for **5**; however, the HRESIMS data revealed that this metabolite possessed one additional degree of unsaturation (*m/z* 247.0573 [M+Na]<sup>+</sup>, calcd for C<sub>11</sub>H<sub>12</sub>O<sub>5</sub>Na 247.0582). The downfield shift of C-5 from δ 55.9 in **5** to δ 87.7 in **6** indicated that the C-7 hydroxyl group formed a new ether linkage to C-5 resulting in a dioxygenated carbon atom. This was readily confirmed with the appearance of two new <sup>3</sup>J<sub>H-C</sub> HMBC correlations extending from H-5 to C-7 (Fig. 2). Compound **6** has been given the trivial name clearanol B. The metabolite degraded upon storage and consequently its relative/absolute configuration was not investigated.

A molecular formula of C<sub>35</sub>H<sub>32</sub>O<sub>14</sub> was established for compound **7** (7 mg)<sup>11</sup> (Fig. 1) based on HRESIMS (*m/z* 699.1685 [M+Na]<sup>+</sup>, calcd for C<sub>35</sub>H<sub>32</sub>O<sub>14</sub>Na 699.1689), which accounted for 20 degrees of unsaturation. Surprisingly, the <sup>13</sup>C NMR data for **7** exhibited only 18 resonances and 17 of the resonances had nearly identical chemical shifts to co-occurring metabolite **1** (C<sub>17</sub>H<sub>16</sub>O<sub>7</sub>). Therefore, we suspected that **7** was a symmetric dimeric compound. After accounting for 17 carbon resonances attributable to **1**, a single new methylene signal remained (C-18) that we deduced would serve as a bridge between two sulochrin moieties. <sup>1</sup>H–<sup>13</sup>C HMBC correlations were observed from the new H-18 methylene protons to C-14/14', C-15/15', and C-16/16', which confirmed that the methylene served as a link between C-15 and C-15'. Compound **7** is structurally similar to acremonidin D, which is the dimerized

product of the demethoxy sulochrin analog, acremonidin E.<sup>12</sup> Metabolite **7** has been assigned the trivial name disulochrin.

The Dothideomycete isolate was grown on Cheerios® supplemented with a 3% sucrose solution and the cultures were extracted with ethyl acetate to afford a brown gummy extract. The extract was subjected to Sephadex LH-20 chromatography and C18 HPLC. This provided five metabolites, two of which were identified by dereplication. Compound **8** (5 mg) was determined to be (R)-4,8-dihydroxy-6-methoxy-4,5-dimethyl-3-methyleneisochroman-1-one<sup>13,14</sup> and **9** (20 mg) was identified as (R)-7-hydroxy-3-((S)-1-hydroxyethyl)-5-methoxy-3,4-dimethylisobenzofuran-1(3H)-one.<sup>14</sup> The absolute configurations of the metabolites were confirmed by comparison of their circular dichroism data to those reported by Tayone et al.<sup>14</sup>

Compound **10** (3 mg)<sup>15</sup> was assigned the molecular formula C<sub>13</sub>H<sub>14</sub>O<sub>4</sub> by HRESIMS (*m/z* 233.0815 [M-H]<sup>-</sup>, calcd for C<sub>13</sub>H<sub>13</sub>O<sub>4</sub>, 233.0814), which accounted for seven degrees of unsaturation. The considerable similarities of the <sup>1</sup>H and <sup>13</sup>C NMR data for **10** and **8** suggested that these metabolites were structurally related. However, a notable difference between the two metabolites concerned C-8 which was shifted upfield from δ 72.1 in **8** to δ 36.0 in **10**. In addition, the <sup>1</sup>H–<sup>13</sup>C HSQC experiment demonstrated that the C-8 quaternary carbon in **8** was a methine in **10**. These data indicated that **10** was the C-8 de-hydroxy analog of metabolite **8**. Subsequent analysis of all the <sup>1</sup>H–<sup>13</sup>C HMBC correlation data led to the assignment of the planar structure of **10** (Fig. 2). We have given compound **10** the name clearanol C.

Compounds **11**<sup>16</sup> and **12**<sup>17</sup> were obtained as a 1:1 mixture (39 mg) of structurally-related metabolites that resisted chromatographic separation by C18 HPLC with aqueous MeOH and acetonitrile. However, examination of the mixture by <sup>1</sup>H and <sup>13</sup>C NMR led us to conclude that the structures of both metabolites could be deduced in their combined form. HRESIMS performed on the mixture yielded a single peak at *m/z* 291.0839 [M+Na]<sup>+</sup>, which represented a molecular formula of C<sub>13</sub>H<sub>16</sub>O<sub>6</sub>, (calcd for C<sub>13</sub>H<sub>16</sub>O<sub>6</sub>Na, 291.0845) for both compounds. In view of the identical molecular formulae and very similar NMR data for these metabolites, it was deduced that **11** and **12** represented a diastereomeric mixture. We also noted that **11** and **12** showed considerable similarity to **9** with the following exceptions: the highfield C-10 methyl doublet protons found in **9** (δ 0.86, *J* = 6.3 Hz) were missing and instead were replaced by downfield diastereotopic methylene double doublets in **11** (δ 3.83, *J* = 11.3, 3.5 Hz and δ 3.53, *J* = 11.3, 8.2 Hz) and **12** (δ 3.34, *J* = 11.3, 8.6 Hz and δ 3.14, *J* = 11.3, 2.7 Hz). This suggested that the additional oxygen atoms in the molecular formulae of **11** and **12** were accounted for by new C-10 hydroxyl groups. <sup>1</sup>H–<sup>13</sup>C HMBC correlations from the C-10 protons of both compounds to their respective C-8 and C-9 carbon atoms (Fig. 2) confirmed these assignments. Further analysis of the <sup>1</sup>H–<sup>13</sup>C HMBC correlation data for both metabolites provided verification of the planar structures of **11** and **12** and we have given these compounds the trivial names clearanols D and E, respectively.

All compounds were tested in bioassays that were performed as previously described.<sup>18,19</sup> Compound **7** exhibited modest activity against methicillin-resistant *Staphylococcus aureus* (ATCC 700787) by completely inhibiting its growth at a concentration of 100 µg/mL; however, growth was not restricted at 10 µg/mL. None of the other compounds inhibited the growth of *S. aureus* or *Klebsiella pneumoniae* (ATCC 51503). Metabolite **10** demonstrated modest growth inhibition effects against polyene-resistant *Candida albicans* (ATCC 38245) and *Aspergillus fumigatus* (FGSC A1100) by reducing growth 61% and 62%, respectively, at 100 µg/mL. None of the other metabolites inhibited the growth of fungi or pancreatic cancer cells (MIA-PaCa-2). Compounds **8** and **10** inhibited *C. albicans* (DAY185) biofilm formation with MIC values of 86 ± 3 and 101 ± 3 µM, respectively. None of the other metabolites

demonstrated activity in the biofilm assay. Based on these studies, we surmise that microbial mat communities offer intriguing potential for encountering unusual fungal taxa with the capacity to generate new bioactive natural products.

## Acknowledgements

Financial support from the National Institutes of Health (1R01GM092219 and 1R01AI085161) is greatly appreciated. The X-ray diffractometer was purchased through a Grant from the National Science Foundation (CHE-0130835).

## Supplementary data

Supplementary data (X-ray diffraction experiment for **5**) associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.tetlet.2012.05.156>.

## References and notes

- Robertson, C. E.; Spear, J. R.; Harris, J. K.; Pace, N. R. *Appl. Environ. Microbiol.* **2009**, *75*, 1801–1810.
- Feazel, L. M.; Spear, J. R.; Berger, A. B.; Harris, J. K.; Frank, D. N.; Ley, R. E.; Pace, N. R. *Appl. Environ. Microbiol.* **2008**, *74*, 329–332.
- Hall-Stoodley, L.; Costerton, J. W.; Stoodley, P. *Nat. Rev. Microbiol.* **2004**, *2*, 95–108.
- Seckbach, J.; Oren, A. *Microbial Mats: Modern and Ancient Microorganisms in Stratified Systems*; Springer: New York, 2010.
- Ley, R. E.; Harris, J. K.; Wilcox, J.; Spear, J. R.; Miller, S. R.; Bebout, B. M.; Maresca, J. A.; Bryant, D. A.; Sogin, M. L.; Pace, N. R. *Appl. Environ. Microbiol.* **2006**, *72*, 3685–3695.
- Phelan, V. V.; Liu, W.-T.; Pogliano, K.; Dorrestein, P. C. *Nat. Chem. Biol.* **2012**, *8*, 26–35.
- Emerson, D.; Fleming, E. J.; McBeth, J. M. *Ann. Rev. Microbiol.* **2010**, *64*, 561–583.
- Clearanol A (5)**: colorless crystals from MeOH; mp 138–140 °C; [α]<sub>D</sub><sup>24</sup> –88 (c 0.4, MeOH); UV (MeOH) λ<sub>max</sub> nm (log ε) 210 (4.35), 290 (3.91); IR (NaCl) ν<sub>max</sub> 3387, 2972, 2346, 1700, 1629, 1560, 1452, 1406, 1246, 1017 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, methanol-*d*<sub>4</sub>) δ 5.75 (1H, d, *J* = 1.5 Hz, H-9a), 5.67 (1H, s, H-3), 5.63 (1H, d, *J* = 1.5 Hz, H-9b), 4.64 (1H, q, *J* = 6.6 Hz, H-7), 4.44 (1H, d, *J* = 11.7 Hz, H-5a), 4.33 (1H, d, *J* = 11.7 Hz, H-5b), 3.94 (3H, s, H-11), 1.30 (3H, d, *J* = 6.6 Hz, H-10); <sup>13</sup>C NMR (125 MHz, methanol-*d*<sub>4</sub>) δ 172.9 (C-4), 166.1 (C-2), 163.1 (C-8a), 146.0 (C-8), 119.8 (C-9), 114.4 (C-4a), 89.5 (C-3), 68.3 (C-7), 57.5 (C-11), 55.9 (C-5), 22.6 (C-10); HRESIMS *m/z* 227.0909 [M+H]<sup>+</sup> (calcd for C<sub>11</sub>H<sub>15</sub>O<sub>5</sub>, 227.0914).
- Cambridge Crystallographic Data Centre no. CCDC 876681; experimental parameters are provided in the online Supplementary data accompanying this report.
- Clearanol B (6)**: amorphous powder; UV (MeOH) λ<sub>max</sub> nm (log ε) 206 (4.29), 298 (3.84); IR (NaCl) ν<sub>max</sub> 3345, 2922, 2853, 2347, 1707, 1609, 1564, 1461, 1408, 1260, 1055 cm<sup>-1</sup>; <sup>1</sup>H NMR (400 MHz, methanol-*d*<sub>4</sub>) δ 5.97 (1H, d, *J* = 1.6 Hz, H-9a), 5.83 (1H, s, H-5), 5.70 (1H, s, H-3), 5.48 (1H, d, *J* = 1.6 Hz, H-9b), 4.92 (1H, *J* = 6.3 Hz, H-7), 3.92 (3H, s, H-11), 1.47 (3H, d, *J* = 6.3 Hz, H-10); <sup>13</sup>C NMR (100 MHz, methanol-*d*<sub>4</sub>) δ 170.7 (C-4), 165.6 (C-2), 153.9 (C-8a), 138.7 (C-8), 114.5 (C-9), 111.4 (C-4a), 90.5 (C-3), 87.7 (C-5), 65.4 (C-7), 57.6 (C-11), 18.4 (C-10); HRESIMS *m/z* 247.0573 [M+Na]<sup>+</sup> (calcd for C<sub>11</sub>H<sub>12</sub>O<sub>5</sub>Na 247.0582).
- Disulochrin (7)**: yellow amorphous powder; UV (MeOH) λ<sub>max</sub> nm (log ε) 210 (4.64), 288 (3.90); IR (NaCl) ν<sub>max</sub> 3301, 2953, 2352, 1710, 1609, 1589, 1441, 1344, 1145, 1064, 1004 cm<sup>-1</sup>; <sup>1</sup>H NMR (400 MHz, methanol-*d*<sub>4</sub>) δ 6.95 (1H, d, *J* = 2.0 Hz, H-5), 6.65 (1H, d, *J* = 2.0 Hz, H-3), 6.00 (1H, s, H-13), 3.98 (1H, s, H-18), 3.70 (3H, s, H-7), 3.62 (3H, s, H-9), 2.17 (3H, s, H-17); <sup>13</sup>C NMR (75 MHz, methanol-*d*<sub>4</sub>) δ 202.4 (C-10,10'), 168.2 (C-8,8'), 162.7 (C-12,12'), 159.8 (C-4,4'), 159.5 (C-16,16'), 158.8 (C-2,2'), 149.3 (C-14,14'), 130.1 (C-6,6'), 128.4 (C-1,1'), 119.0 (C-15,15'), 110.8 (C-11,11'), 109.4 (C-13,13'), 109.0 (C-5,5'), 104.4 (C-3,3'), 56.7 (C-7,7'), 52.7 (C-9,9'), 22.0 (C-18), 21.1 (C-17,17'); HRESIMS *m/z* 699.1685 [M+Na]<sup>+</sup> (calcd for C<sub>35</sub>H<sub>32</sub>O<sub>14</sub>Na 699.1689).
- He, H.; Bigelis, R.; Solum, E. H.; Greenstein, M.; Carter, G. T. *J. Antibiot.* **2003**, *56*, 923–930.
- Chinworrungsee, M.; Kittakoop, P.; Isaka, M.; Chanphen, R.; Tanticharoen, M.; Thebtaranonth, Y. *J. Chem. Soc., Perkin Trans. 1* **2002**, 2473–2476.
- Tayone, W. C.; Honma, M.; Kanamaru, S.; Noguchi, S.; Tanaka, K.; Nehira, T.; Hashimoto, M. *J. Nat. Prod.* **2011**, *74*, 425–429.
- Clearanol C (10)**: amorphous solid; [α]<sub>D</sub><sup>22</sup> +39 (c 0.06, MeOH); UV (MeOH) λ<sub>max</sub> nm (log ε) 220 (3.83) 270 (3.47), 312 (3.20); IR (NaCl) ν<sub>max</sub> 3623, 2949, 2851, 1651, 1018 cm<sup>-1</sup>; <sup>1</sup>H NMR (400 MHz, methanol-*d*<sub>4</sub>) δ 6.46 (1H, s, H-4), 4.74 (1H, s, H-10a), 4.68 (1H, s, H-10b), 3.99 (1H, q, *J* = 7.0 Hz, H-8), 3.88 (3H, s, H-13), 2.09 (3H, s, H-12), 1.34 (3H, d, *J* = 7.0 Hz, H-11); <sup>13</sup>C NMR (100 MHz, methanol-*d*<sub>4</sub>) δ 168.4 (C-5), 166.6 (C-1), 164.4 (C-3), 159.1 (C-9), 143.8 (C-7), 115.5 (C-6), 99.5 (C-2), 98.5 (C-4), 96.2 (C-10), 56.6 (C-13), 36.0 (C-8), 22.7 (C-11), 10.2 (C-12); HRESIMS *m/z* 233.0815 [M-H]<sup>-</sup> (calcd for C<sub>13</sub>H<sub>13</sub>O<sub>4</sub>, 233.0814).

16. *Clearanol D* (**11**): amorphous solid (1:1 mixture with **12**); UV (MeOH)  $\lambda_{\text{max}}$  nm (log  $\epsilon$ ) 218 (4.22), 260 (3.87), 300 (3.59); IR (NaCl)  $\nu_{\text{max}}$  3596, 3089, 1737, 1651, 1472, 1373, 1319, 1247, 1220, 1153, 1067  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (400 MHz, methanol- $d_4$ )  $\delta$  6.47 (1H, s, H-4), 4.13 (1H, dd,  $J = 3.7$  and 8.0 Hz, H-9), 3.87 (3H, s, H-13), 3.83 (1H, dd,  $J = 3.5$  and 11.3 Hz, H-10a), 3.53 (1H, dd,  $J = 8.2$  and 11.3 Hz, H-10b), 2.18 (3H, s, H-12), 1.67 (3H, s, H-11);  $^{13}\text{C}$  NMR (100 MHz, methanol- $d_4$ )  $\delta$  171.9 (C-1), 166.4 (C-5), 157.8 (C-3), 153.1 (C-7), 113.4 (C-6), 105.4 (C-2), 99.5 (C-4), 90.6 (C-8), 75.7 (C-9), 63.9 (C-10), 56.9 (C-13), 22.08 (C-11), 11.4 (C-12); HRESIMS  $m/z$  291.0839  $[\text{M}+\text{Na}]^+$  (calcd for  $\text{C}_{13}\text{H}_{16}\text{O}_6\text{Na}$ , 291.0845).
17. *Clearanol E* (**12**): amorphous solid (1:1 mixture with **11**); UV (MeOH)  $\lambda_{\text{max}}$  nm (log  $\epsilon$ ) 218 (4.22), 260 (3.87), 300 (3.59); IR (NaCl)  $\nu_{\text{max}}$  3596, 3089, 1737, 1651, 1472, 1373, 1319, 1247, 1220, 1153, 1067  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (400 MHz, methanol- $d_4$ )  $\delta$  6.49 (1H, s, H-4), 4.11 (1H, dd,  $J = 2.7$  and 8.2 Hz, H-9), 3.88 (3H, s, H-13), 3.34 (1H, dd,  $J = 8.6$  and 11.3 Hz, H-10a), 3.14 (1H, dd,  $J = 2.7$  and 11.3 Hz, H-10b), 2.15 (3H, s, H-12), 1.73 (3H, s, H-11);  $^{13}\text{C}$  NMR (100 MHz, methanol- $d_4$ )  $\delta$  171.5 (C-1), 166.6 (C-5), 158.3 (C-3), 152.4 (C-7), 113.3 (C-6), 104.6 (C-2), 99.9 (C-4), 90.4 (C-8), 76.2 (C-9), 63.3 (C-10), 56.9 (C-13), 22.11 (C-11), 11.7 (C-12); HRESIMS  $m/z$  291.0839  $[\text{M}+\text{Na}]^+$  (calcd for  $\text{C}_{13}\text{H}_{16}\text{O}_6\text{Na}$ , 291.0845).
18. Wang, X.; You, J.; King, J. B.; Powell, D. R.; Cichewicz, R. H. *J. Nat. Prod.* **2012**, *75*, 707–715.
19. Henrikson, J. C.; Ellis, T. K.; King, J. B.; Cichewicz, R. H. *J. Nat. Prod.* **2011**, *74*, 1959–1964.